



Assisted dying and non-assisted suicide



Non-assisted suicides in Oregon have risen by one third, despite legalising assisted dying ^{1, 2}

In 2021, only two patients were referred for psychiatric evaluation ³



References

- Oregon Health Authority Public Health Division. <https://www.oregon.gov/oha/PH/Dis easesConditions/InjuryFatalityData/P ages/nvdrs.aspx>
- CDC suicide mortality: <https://www.cdc.gov/nchs/pressroo m/sosmap/suicide-mortality/suicide.htm>
- Oregon Death with Dignity Act: annual reports. 2021.
- Jones DA, et al. *Journal of Ethics in Mental Health*, 2022; 11:1.
- Centre for Suicide Prevention <https://www.suicideinfo.ca/resource/ suicide-stats-canada-provinces/>
- Suicides among people diagnosed with severe health conditions, England: 2017 to 2020. London: ONS, 2022
- Rabkin J et al. *Neurology*, 2016; **87**(13): 1320-8.
- Hartog ID et al. *BMC Geriatrics*, 2020; **20**: 342-356.
- Kim SYH, et al *JAMA Psychiatry*. 2016; **73**(4): 362-368
- Appelbaum PS. *JAMA Psychiatry*, 2016; **73**(4): 325-6.



In Europe suicide rates have not decreased after the legalisation of assisted dying.⁴

Dutch non-assisted suicides have increased compared with Germany.⁴

Belgium now has the highest non-assisted suicide rate in women in Europe.⁴



In Canada non-assisted suicide rates increased 2016-19 (11-12.2/100k) then reduced in 2020 (10.1/100k) ⁵



In UK data confirms previous work that the risk of suicide is highest in the year after diagnosis, then the risk reduces.⁶ Suicide prevention is key.

In progressive neurological disease, a wish to die is strongly related to depression ⁷

Three quarters of those with a wish to die report being lonely and 60% have a clinical depression.⁸

Of Dutch psychiatric assisted deaths, 55% were depressed, 11% had no independent psychiatric input, and 20% never had a psychiatric inpatient stay ^{9, 10}

